



Learn what it can mean for you

The **Hello**CUVITRU Free Trial Program was designed with you in mind

It's about making the first step toward weekly to every-other-week infusions. With the HelloCUVITRU Free Trial Program, you can try CUVITRU for free to determine, along with your doctor, if it's right for you.*

*To be eligible, you must be a patient with an ICD-10-verified diagnosis, be a new patient not currently using CUVITRU, and have not been previously enrolled in the HelloCUVITRU program.

ICD-10=International Statistical Classification of Diseases, Tenth Revision.



Highlights of the HelloCUVITRU Free Trial Program

- ✓ The entire cost of your first 4 CUVITRU infusions are free, including the pump, supplies, and administration training. Everything you need will be shipped directly to your home.
- ✓ A nurse will help administer your CUVITRU therapy and can work with you to optimize your therapy administration parameters.
- ✓ Together, you can customize the number of infusion sites, the volume infused per site, and how fast and how often you infuse.

What is CUVITRU?

CUVITRU is a ready-to-use liquid medicine that is given under the skin (subcutaneously) to treat primary immunodeficiency (PI) in people 2 years and older.

IMPORTANT SAFETY INFORMATION

What is the most important information I need to know about CUVITRU?

CUVITRU can cause the following serious reactions:

- Severe allergic reactions causing difficulty in breathing or skin rashes
- Decreased kidney function or kidney failure
- Blood clots in the heart, brain, lungs, or elsewhere in the body
- Severe headache, drowsiness, fever, painful eye movements, or nausea and vomiting
- Dark colored urine, swelling, fatigue, or difficulty breathing

Please see additional safety information throughout and click for [Information For Patients](#) and discuss with your HCP.

The **Hello**CUVITRU process



Take the first step

Once you and your HCP decide to enroll you in the **HelloCUVITRU Free Trial Program**, your HCP will verify your eligibility* and start the enrollment process.



What you need to do

If you do not have epinephrine or antihistamines at home, you will need to ask your HCP whether you should have rescue medications, such as antihistamines or epinephrine. As a safety measure, epinephrine or antihistamines need to be on hand for your first free trial CUVITRU infusion.

Meet your Case Manager

Before your first infusion, you'll receive a welcome call from a HelloCUVITRU Case Manager who will help you with the following:

- Completing your HelloCUVITRU Free Trial Form if needed.
- Connecting with RxCrossroads, the SP that will coordinate the shipment of your medication.
- Connecting with a nurse who will help schedule your in-home administration training.



What you need to do

Keep an eye out for a call or voicemail from your Case Manager (this call may come from a number you do not recognize).

Treat at home with support

A nurse will come to your home to administer your CUVITRU infusions and provide initial administration training. This will include:

- How to set up the supplies for infusion of CUVITRU.
- Detailed instructions for administration.



What you need to do

Have your HelloCUVITRU Checklist ready and make sure your nurse fills in the appropriate information.

*To be eligible, you must be a patient with an ICD-10–verified diagnosis, be a new patient not currently using CUVITRU, and have not been previously enrolled in the HelloCUVITRU program.

HCP=healthcare professional; SP=specialty pharmacy.

Please see safety information, including Warning about Blood Clots, throughout and click for [Information For Patients and discuss with your HCP.](#)



cuvitru

[Immune Globulin Subcutaneous (Human)] 20%

Infusions tailored to you^{†‡}



Fewer needlesticks

In one study, most participants (84.9%) experienced 1 to 2 needlesticks.



Faster infusions

CUVITRU can be infused at the fastest rates and highest volumes with the fewest infusions of any subcutaneous immune globulin.[‡]



Flexible schedule

CUVITRU gives you and your HCP the freedom to tailor your PI treatment to your lifestyle with weekly or every-other-week infusions.

[†]CUVITRU was studied in 77 people with PI ≥ 2 years of age. The average infusion time was 0.95 hours (range, 0.2-6.4 hours).

[‡]You'll infuse your first 2 infusions at 10 to 20 mL/hour/site. After that, you'll be able to increase your rate up to 60 mL/hour/site, as tolerated. Infuse at up to 4 sites simultaneously.

PI=primary immunodeficiency.



HelloCUVITRU Checklist

This checklist can be used by you and your nurse to help monitor your infusions during the free trial program.

IMPORTANT SAFETY INFORMATION (Continued)

Who should not use CUVITRU?

Do not use CUVITRU if you:

- Have had a severe allergic reaction to immune globulin or other blood products.
- Have a condition called selective (or severe) immunoglobulin A (IgA) deficiency.

What should I avoid while taking CUVITRU?

- CUVITRU can make vaccines (like measles/mumps/rubella or chickenpox vaccines) not work as well for you. Before you get any vaccines, tell your healthcare provider (HCP) that you take CUVITRU.
- Tell your HCP if you are pregnant, or plan to become pregnant, or if you are nursing.

What are the possible or reasonably likely side effects of CUVITRU?

CUVITRU can cause serious side effects. If any of the following problems occur after starting CUVITRU, stop the infusion immediately and contact your HCP or call emergency services:

- Hives, swelling in the mouth or throat, itching, trouble breathing, wheezing, fainting or dizziness. These could be signs of a serious allergic reaction.
- Bad headache with nausea, vomiting, stiff neck, fever, and sensitivity to light. These could be signs of irritation and swelling of the lining around your brain.
- Reduced urination, sudden weight gain, or swelling in your legs. These could be signs of a kidney problem.
- Pain, swelling, warmth, redness, or a lump in your legs or arms. These could be signs of a blood clot.
- Brown or red urine, fast heart rate, yellow skin or eyes. These could be signs of a liver or blood problem.
- Chest pain or trouble breathing, or blue lips or extremities. These could be signs of a serious heart or lung problem.
- Fever over 100°F. This could be sign of an infection.

Please see additional safety information, including Warning about Blood Clots, throughout and click for Information For Patients and discuss with your HCP.

HelloCUVITRU

After your free trial, if you and your HCP decide CUVITRU is right for you, you will need a prescription. The insurance approval process can be lengthy. **Reach out to your HCP's office prior to your last infusion to allow enough time for this process.**

You will also work with a different SP to manage your new prescription. As part of that onboarding process, your SP may provide you with a new nurse to help you formally onboard onto CUVITRU as ongoing therapy.

[Click here](#) to learn more about patient experiences with CUVITRU.



If you're eligible, the HelloCUVITRU program lets you try CUVITRU for free to see if it's right for you*

For additional information or to learn how to enroll in the HelloCUVITRU Free Trial Program, contact your HCP, visit **[CUVITRU.com](https://www.cuvitru.com)**, or call **1-866-861-1750**.

*To be eligible, you must be a patient with an ICD-10-verified diagnosis, be a new patient not currently using CUVITRU, and have not been previously enrolled in the HelloCUVITRU program.

IMPORTANT SAFETY INFORMATION (Continued)

What are the possible or reasonably likely side effects of CUVITRU? (Continued)

The following one or more possible side effects may occur at the site of infusion. These generally go away within a few hours, and are less likely after the first few infusions.

- Mild or moderate pain
- Redness
- Itching

The most common side effects that may occur are:

- Headache
- Nausea
- Fatigue
- Diarrhea
- Vomiting

These are not all the possible side effects. Talk to your HCP about any side effect that bothers you or that does not go away.

For additional safety information, including Warning about Blood Clots, click for [Information For Patients](#) and discuss with your HCP.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.